Mucovent

Tablets

Composition:
Each Tablet contains:

Salbutamol (as sulphate)	1.0 mg.
Bromhexine HCl	4.0 mg.

Properties:
The product contains two effective recent respiratory therapeutics intended to clear the respiratory airways and to improve both expiratory volume and vital capacity in bronchitic patients.
Salbutamol sympathomimetic bronchodilator acting specifically and Selectively on Beta-2 sympathetic receptors of the bronchial tree by so breaking down bronchospasm.
Bromhexine the recent mucokinetic agent which is claimed to act directly on bronchial glands increasing both water and solid content of bronchial mucous and depolymerizes muco polysaccharides in the mucous resulting in reduction of viscosity and facilitales expectoration of bronchial secretions .
There by the products cuts the viscous circle of cough (irritation – cough -spasm) in two sites and also provides airways.

Indications and usage :
Cough of lower respiratory tract origin associated with bronchospasm e.g. asthma, asthmatic bronchitis, emphysema, bronchiectasis, iungabcess, external irritant. Smokers cough etc.

Contra –Indications:
1-Significant cardiac arrhythmia.
2-Peptic ulceration

Side effects:
Very rare and unlike to be happened mainly due to salbutamol content:
-Fine tremors of hand and fingers.
-Headache, dizziness, tachycardia, insomnia.
Due to both ingredients very uncommon upper GIT troubles, nausea, vomiting

Drug interactions:
Due to salbutamol:
1-increase the effects of
a-other adrenergics.
b-Xanthines (C.N.S. stimulation) .
2-if taken with beta – adrenergic blocking actions of both may be inhibited.
3-Tricyclic antidepressants may increases the cardiac effects of Salbutamol.

Warnings and precautions:
-Cardiac disorders.
-Vascular disorders.
-Diabetes mellitus and thyrotoxicosis
If taken with sympathomimetics or MAOI.
Note:
Drug tolerance may be developed with prolonged use of salbutamol. Effectiveness of the drug is regained by discontinuing for short period.

Dosage and administration:
Adults : 2 tablets 3-4 times daily.
Children below 12 years 1 tablet 3 times daily.

Packing :
Boxes of 20 tablets.

- Medicament is a product which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine. it's benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.
- Keep medicament out of children reach.

Council of Arab Health Ministers and Union of Arab Pharmacists

MISR COMPANY FOR PHARMACEUTICALS